

Gröbner Basis and Planar Graph

Dang Vu Giang

Hanoi Institute of Mathematics

18 Hoang Quoc Viet, 10307 Hanoi, Vietnam

e-mail: <dangvugiang@yahoo.com>

February 27, 2012

Abstract. Using the Gröbner basis of an ideal generated by a family of polynomials we prove that every planar graph is 4-colorable.

Keywords:

AMS subject classification: 13P10

We are given a graph G with n vertices. Between two any vertices there is at most one edge. We want to color the vertices by 4 colors such that no two vertices connected by an edge are colored the same way. We let $\zeta = i \in \mathbb{C}$ be the 4th primitive root of unity ($i^4 = 1$). Let x_1, x_2, \dots, x_n be variables representing the distinct vertices of the graph G . Each vertex is to be assigned one of colors $1, i, i^2, i^3$. This means that $x_k^4 - 1 = 0$ for $k = 1, 2, \dots, n$. Moreover, if x_j and x_k are connected then $x_j^3 + x_j^2 x_k + x_j x_k^2 + x_k^3 = 0$. Let I be the ideal generated by $x_k^4 - 1$ for $k = 1, 2, \dots, n$ and $x_j^3 + x_j^2 x_k + x_j x_k^2 + x_k^3$ if x_j and x_k are connected. We should prove that $I \neq \mathbb{C}[z_1, z_2, \dots, z_n]$.

Acknowledgement. Deepest appreciation is extended towards the NAFOSTED (the National Foundation for Science and Technology Development in Vietnam) for the financial support.

References

- [1] W. Adams and P. Loustau, An introduction to Gröbner basis, AMS 1996.
- [2]